

CONSTRUCTING r -MATRICES ON SIMPLE LIE SUPERALGEBRAS

GIZEM KARAALI

ABSTRACT. We construct r -matrices for simple Lie superalgebras with non-degenerate Killing forms using Belavin-Drinfeld type triples. This construction gives us the standard r -matrices and some nonstandard ones.

1. INTRODUCTION

Let $\mathfrak{g}$ be a Lie algebra with a non-degenerate $\mathfrak{g}$ -invariant bilinear form $(\cdot, \cdot)$. Denote by Ω the element of $(\mathfrak{g} \otimes \mathfrak{g})^{\mathfrak{g}}$ that corresponds to the quadratic Casimir element in $\mathfrak{U}\mathfrak{g}$ of $\mathfrak{g}$. Then the **classical Yang-Baxter equation (CYBE)** for an element $r \in \mathfrak{g} \otimes \mathfrak{g}$ is:

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0.$$

A solution r to the classical Yang-Baxter equation is called a **classical r -matrix** (or simply an r -**matrix**). An r -matrix r is called **non-degenerate** if it satisfies:

$$r^{12} + r^{21} = \Omega.$$

In [1] and [2] Belavin and Drinfeld classified such r -matrices. The solutions in this classification are parametrized by triples $(\Gamma_1, \Gamma_2, \tau)$ (called **admissible triples**) where the Γ_i are certain subsets of the set Γ of simple roots of $\mathfrak{g}$, and $\tau : \Gamma_1 \rightarrow \Gamma_2$ is an isometric bijection. They proved that for each admissible triple and some fixed $r_0 \in \mathfrak{g} \otimes \mathfrak{g}$ there exists a unique non-degenerate classical r -matrix, and conversely that each non-degenerate classical r -matrix can be associated with such data.

In this paper we construct classical r -matrices using analogs of the Belavin-Drinfeld data for simple Lie superalgebras with non-degenerate Killing form. We first start with a review of the situation in the Lie algebra case. Thus in Section 2 we give an overview of the Belavin-Drinfeld result for simple Lie algebras. Next, in Section 3, we recall some basic definitions and results about simple Lie superalgebras. Then after developing the necessary ingredients we state our main theorem. The next three sections of the paper are devoted to the proof of this theorem. Then in Section 7 we construct various r -matrices for the Lie superalgebra $sl(2, 1)$ using the main theorem.

This theorem is very much in the spirit of the Belavin-Drinfeld result. It tells us that, given a Belavin-Drinfeld type triple, one can construct a non-degenerate r -matrix in a way similar to the construction in the Lie algebra case. One should point out, however, that this is not a complete classification result. The theorem gives us r -matrices, but does not tell us whether we can always extract a Belavin-Drinfeld type triple from a given r -matrix. We discuss this briefly in the last section of the paper.

Acknowledgments. I would like to thank Nicolai Reshetikhin for fruitful discussions, Vera Serganova for her close attention and useful suggestions, and Milen Yakimov for introducing me to the Belavin-Drinfeld result. This work was supported by the NSF grant DMS-0070931.

2. CLASSIFICATION THEOREM FOR LIE ALGEBRAS

Here we recall briefly the main result of [1] and [2] for Lie algebras. Let $\mathfrak{g}$ be a simple Lie algebra. Fixing a positive Borel subalgebra $\mathfrak{b}_+$ determines a Cartan subalgebra $\mathfrak{h}$ for $\mathfrak{g}$, and we can talk about positive roots, simple roots etc. Hence we can define $\Gamma = \{\alpha_1, \alpha_2, \dots, \alpha_r\}$ to be the set of all simple roots of $\mathfrak{g}$. We will be interested in **admissible triples**, i.e. triples $(\Gamma_1, \Gamma_2, \tau)$ where $\Gamma_i \subset \Gamma$ and $\tau : \Gamma_1 \rightarrow \Gamma_2$ is a bijection such that

- (1) for any $\alpha, \beta \in \Gamma_1$, $(\tau(\alpha), \tau(\beta)) = (\alpha, \beta)$;
- (2) for any $\alpha \in \Gamma_1$ there exists a $k \in \mathbb{N}$ such that $\tau^k(\alpha) \notin \Gamma_1$.¹

We will also need a continuous parameter r_0 , an element of $\mathfrak{h} \otimes \mathfrak{h}$ which satisfies the following equations:

$$r_0^{12} + r_0^{21} = \Omega_0 \quad (1)$$

$$(\tau(\alpha) \otimes 1)(r_0) + (1 \otimes \alpha)(r_0) = 0 \text{ for all } \alpha \in \Gamma_1 \quad (2)$$

where $\Omega_0 \in \mathfrak{h} \otimes \mathfrak{h}$ is the $\mathfrak{h}$ -component of the quadratic Casimir element Ω of $\mathfrak{g}$.

Fix a system of Weyl-Chevalley generators $X_\alpha, Y_\alpha, H_\alpha$ for $\alpha \in \Gamma$. Recall that these elements generate the Lie algebra $\mathfrak{g}$ with the defining relations: $[X_{\alpha_i}, Y_{\alpha_j}] = \delta_{i,j} H_{\alpha_j}$, $[H_{\alpha_i}, X_{\alpha_j}] = a_{i,j} X_{\alpha_j}$ and $[H_{\alpha_i}, Y_{\alpha_j}] = -a_{i,j} Y_{\alpha_j}$ for all $\alpha_i, \alpha_j \in \Gamma$, where $a_{i,j} = \alpha_j(H_{\alpha_i}) = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)}$, along with the well-known Serre relations.

Denote by $\mathfrak{g}_i$ the subalgebra of $\mathfrak{g}$ generated by the elements $X_\alpha, Y_\alpha, H_\alpha$ for all $\alpha \in \Gamma_i$. We define a map φ by:

$$\varphi(X_\alpha) = X_{\tau(\alpha)} \quad \varphi(Y_\alpha) = Y_{\tau(\alpha)} \quad \varphi(H_\alpha) = H_{\tau(\alpha)}$$

for all $\alpha \in \Gamma_1$. Then this can be extended to an isomorphism $\varphi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ because the relations between $X_\alpha, Y_\alpha, H_\alpha$ for $\alpha \in \Gamma_1$ will be the same as the relations between $X_{\tau(\alpha)}, Y_{\tau(\alpha)}, H_{\tau(\alpha)}$ for $\alpha \in \Gamma_1$. Note that this requires the first property of τ , namely that it is an isometry. Next extend τ to a bijection between the $\overline{\Gamma}_i$, where $\overline{\Gamma}_i$ is the set of those roots which can be written as a nonnegative integral linear combination of the elements of Γ_i . In each root space $\mathfrak{g}_\alpha$, we choose an element e_α such that $(e_\alpha, e_{-\alpha}) = 1$ for any α and $\varphi(e_\alpha) = e_{\tau(\alpha)}$ for all $\alpha \in \overline{\Gamma}_1$.² Finally we define a partial order on the set of all positive roots:

$$\alpha < \beta \text{ if and only if there exists a } k \in \mathbb{N} \text{ such that } \beta = \tau^k(\alpha)$$

¹The expression $\tau^k(\alpha)$ has a meaning only if the expressions $\tau^j(\alpha)$ for all $j < k$ are elements of Γ_1 . So this condition actually may be translated as saying that $\tau^k(\alpha)$ does not make sense for large enough k .

²Such e_α can always be chosen consistently if “there are no cycles”, i.e. if τ satisfies the second property. Otherwise, if there is a cycle of simple roots $\alpha_1, \dots, \alpha_k$ such that $\tau(\alpha_i) = \alpha_{i+1}$ for $i = 1, 2, \dots, k-1$ and $\tau(\alpha_k) = \alpha_1$ and τ is an isometry, then we must have that $(\alpha_i, \alpha_{i+1}) = 0$ for all $i = 1, 2, \dots, k-1$, since no cycles are allowed in Dynkin diagrams. Then if s is the smallest integer such that $(\alpha_1, \alpha_{1+s}) \neq 0$, then $s > 1$ and $s|k$. So our cycle must have s disconnected subgraphs of length k/s . Then we can still choose e_α consistently, provided we allow only cycles as above.

Note that in this case we will have $\alpha \in \overline{\Gamma}_1, \beta \in \overline{\Gamma}_2$. Clearly one needs property 2 for a partial order; no cycles are allowed in partial orders.

Now we can state the Belavin-Drinfeld theorem:

Theorem 1. (1) Let $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ satisfy Equations 1 and 2. Then the element r of $\mathfrak{g} \otimes \mathfrak{g}$ defined by:

$$r = r_0 + \sum_{\alpha > 0} e_{-\alpha} \otimes e_{\alpha} + \sum_{\alpha, \beta > 0, \alpha < \beta} (e_{-\alpha} \otimes e_{\beta} - e_{\beta} \otimes e_{-\alpha})$$

is a solution to the system:

$$r^{12} + r^{21} = \Omega \quad (3)$$

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0 \quad (4)$$

(2) Any solution (up to isomorphism) to the above system can be obtained as above by a suitable choice of an admissible triple $(\Gamma_1, \Gamma_2, \tau)$ and some $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ that satisfies Equations 1 and 2.

The proof of this theorem provided in [2] is actually quite clear; one can also look at [3] for another exposition.

3. THE CONSTRUCTION THEOREM FOR LIE SUPERALGEBRAS

Now our aim is to develop a similar theory for super structures. Let $\mathfrak{g}$ be a simple Lie superalgebra with a non-degenerate Killing form ³.

3.1. The Quadratic Casimir Element: Let $\{I_{\alpha}\}$ be a homogeneous basis for $\mathfrak{g}$ and denote by $\{I_{\alpha}^*\}$ the dual basis of $\mathfrak{g}$ with respect to the non-degenerate (Killing) form. Thus we have:

$$(I_{\alpha}, I_{\beta}^*) = \delta_{\alpha\beta}$$

If we denote the parity of a homogeneous element $x \in \mathfrak{g}$ by $|x|$, then we also have that

$$|I_{\alpha}| = |I_{\alpha}^*|$$

since the supertrace form is consistent [Recall that a bilinear form $(\cdot, \cdot)$ is **consistent** if for any homogeneous $x, y \in \mathfrak{g}$ of different parities, $(x, y) = 0$].

We can write the quadratic Casimir element of $\mathfrak{g}$ as follows:

$$\Omega = \sum_{\alpha} (-1)^{|I_{\alpha}||I_{\alpha}^*|} I_{\alpha} \otimes I_{\alpha}^* = \sum_{\alpha} (-1)^{|I_{\alpha}|} I_{\alpha} \otimes I_{\alpha}^*$$

Example 1. For the special case $\mathfrak{g} = gl(m, n)$ with basis $\{e_{i,j} | 1 \leq i, j \leq m+n\}$, we use the supertrace form to find the dual basis:

$$e_{i,j}^* = (-1)^{[i]} e_{j,i}$$

where

$$[j] = \begin{cases} 0 & \text{if } j \leq m \\ 1 & \text{if } j > m \end{cases}$$

Then this gives us:

$$\Omega = \sum_{\alpha} (-1)^{|I_{\alpha}|} I_{\alpha} \otimes I_{\alpha}^* = \sum_{i,j} (-1)^{|e_{i,j}|} e_{i,j} \otimes (-1)^{[i]} e_{j,i} = \sum_{i,j} (-1)^{[j]} e_{i,j} \otimes e_{j,i}$$

³This implies that $\mathfrak{g}$ is isomorphic to a simple Lie algebra or to one of the following classical Lie superalgebras: $A(m, n)$ with $m \neq n$, $B(m, n)$, $C(n)$, $D(m, n)$ with $m - n \neq 1$, $F(4)$, or $G(3)$. See [4] for details.

3.2. Borel subsuperalgebras and Dynkin diagrams: Let $\mathfrak{h}$ be a Cartan subalgebra. By definition, $\mathfrak{h} \subset \mathfrak{g}_{\overline{0}}$ is a Cartan subalgebra of the even part of our Lie superalgebra. Let $\Delta = \Delta_{\overline{0}} + \Delta_{\overline{1}}$ be the set of all roots of $\mathfrak{g}$ associated with the Cartan subalgebra $\mathfrak{h}$ ⁴, where $\Delta_{\overline{0}}$ and $\Delta_{\overline{1}}$ are the even and odd roots respectively. Fix a Borel subsuperalgebra $\mathfrak{b}$ containing $\mathfrak{h}$. Recall that a Lie subsuperalgebra $\mathfrak{b}$ of a Lie superalgebra $\mathfrak{g}$ is a **Borel subsuperalgebra** if there is some Cartan subsuperalgebra $\mathfrak{h}$ of $\mathfrak{g}$ and some base Γ for Δ , such that

$$\mathfrak{b} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_{\alpha}$$

where Δ^+ consists of all nonnegative integral combinations of the elements of Γ that are in Δ .

In the Lie algebra case subalgebras given by this definition turn out to be maximally solvable, and all maximally solvable subalgebras of a simple Lie algebra are of this type. Thus this definition agrees with the usual definition of a Borel subalgebra as a maximally solvable subalgebra. However Borel subsuperalgebras as defined above are not necessarily maximally solvable.

Example 2. Let $\mathfrak{g}$ be a Lie superalgebra, fix some set Δ_+ of positive roots, and let α be a positive isotropic root. Define $\mathfrak{b}$ as the sum of all the positive root spaces. Then $\mathfrak{b}$ is a Borel subsuperalgebra but is not maximally solvable. The (parabolic⁵) subsuperalgebra $\mathfrak{p} = \mathfrak{b} \oplus \mathfrak{g}_{-\alpha}$ is also solvable.

In fact maximally solvable subsuperalgebras may be even more complicated. [See [6] for a study of maximally solvable subsuperalgebras of $gl(m, n)$ and $sl(m, n)$.] Therefore we choose to define Borels as above instead of using the more traditional characterization by maximal solvability.

Thus Borel subsuperalgebras determine simple roots, and different Borel subsuperalgebras may correspond to different Dynkin diagrams and Cartan matrices. Let us then fix some Borel subsuperalgebra $\mathfrak{b}$, or equivalently some set Γ of simple roots, and the associated Dynkin diagram D . Note that Γ may contain even and odd roots. Another significant difference from the theory of Lie algebras is to be noted here; two Dynkin diagrams of a given Lie superalgebra are not necessarily isomorphic, but can be obtained from one another via a chain of odd reflections.

3.3. The Data for the Theorem: In this setup, let Δ^+ (resp. Δ^-) be the set of all positive (resp. negative) roots, with respect to the chosen $\Gamma = \{\alpha_1, \alpha_2, \dots, \alpha_r\}$.

Now let $\Gamma_1, \Gamma_2 \subset \Gamma$ be two subsets, and $\tau : \Gamma_1 \rightarrow \Gamma_2$ be a bijection. We will say that the triple $(\Gamma_1, \Gamma_2, \tau)$ is **admissible** if:

- (1) for any $\alpha, \beta \in \Gamma_1$, $(\tau(\alpha), \tau(\beta)) = (\alpha, \beta)$;
- (2) for any $\alpha \in \Gamma_1$ there exists a $k \in \mathbb{N}$ such that $\tau^k(\alpha) \notin \Gamma_1$;
- (3) τ preserves grading, i.e. it maps even roots to even ones, and odd roots to odd ones.

For a given admissible triple $(\Gamma_1, \Gamma_2, \tau)$, we define $\overline{\Gamma}_i$ for $i = 1, 2$ as in the Lie algebra case: $\overline{\Gamma}_i$ is the set of those roots which can be written as a nonnegative

⁴In fact Δ is independent of the choice of $\mathfrak{h}$

⁵As in the Lie algebra case, a Lie subsuperalgebra $\mathfrak{p}$ is a **parabolic subsuperalgebra** if $\mathfrak{p}$ contains a Borel.

integral linear combination of the elements of Γ_i . Then we can extend τ linearly to a bijection $\bar{\tau}: \bar{\Gamma}_1 \rightarrow \bar{\Gamma}_2$. Using $\bar{\tau}$, we define a partial order on Δ^+ :

$$\alpha < \beta \text{ if and only if there exists a } k \in \mathbb{N} \text{ such that } \beta = \bar{\tau}^k(\alpha)$$

For any $\alpha \in \Gamma$, pick a nonzero $e_\alpha \in \mathfrak{g}_\alpha$. Then since each of the $\mathfrak{g}_\alpha$ are one dimensional, and the Killing form is a non-degenerate pairing of $\mathfrak{g}_\alpha$ with $\mathfrak{g}_{-\alpha}$ ⁶, we can uniquely pick $e_{-\alpha} \in \mathfrak{g}_{-\alpha}$ such that:

$$(e_\alpha, e_{-\alpha}) = 1.$$

Then we will have, for each $\alpha \in \Gamma$:

$$[e_\alpha, e_{-\alpha}] = (e_\alpha, e_{-\alpha})h_\alpha$$

where h_α is the nonzero vector defined by $(h_\alpha, h) = \alpha(h)$ for all $h \in \mathfrak{h}$. The set $\{h_\alpha | \alpha \in \Gamma\}$ is a basis for $\mathfrak{h}$. Hence we can write Ω_0 , the $\mathfrak{h}$ -part of the quadratic Casimir Ω of $\mathfrak{g}$, as follows:

$$\Omega_0 = \sum_{i=1}^r h_{\alpha_i} \otimes h_{\alpha_i}^*$$

where the set $\{h_\alpha^* | \alpha \in \Gamma\}$ is the basis in $\mathfrak{h}$ dual to $\{h_\alpha | \alpha \in \Gamma\}$

Actually we can choose a nonzero $e_\alpha \in \mathfrak{g}_\alpha$ for each $\alpha \in \Delta$ such that $(e_\alpha, e_{-\alpha}) = 1$ whenever α is positive. Then we compute the duals with respect to the standard (Killing) form:

$$\begin{aligned} e_\alpha^* &= e_{-\alpha} \\ e_{-\alpha}^* &= (-1)^{|\alpha|} e_\alpha \end{aligned}$$

for all positive roots α . Here $|\alpha|$ is the parity of the root α .

Then we can see that the quadratic Casimir element of our Lie superalgebra $\mathfrak{g}$ will be:

$$\begin{aligned} \Omega &= \sum_i (-1)^{|I_i|} I_i \otimes I_i^* \\ &= \sum_{i=1}^r h_{\alpha_i} \otimes h_{\alpha_i}^* + \sum_{\alpha \in \Delta} (-1)^{|e_\alpha|} e_\alpha \otimes e_\alpha^* \\ &= \Omega_0 + \sum_{\alpha \in \Delta^+} (-1)^{|\alpha|} e_\alpha \otimes e_{-\alpha} + \sum_{\alpha \in \Delta^+} e_{-\alpha} \otimes e_\alpha \end{aligned}$$

Example 1 continued: *Let us consider the special case $\mathfrak{g} = gl(m, n)$ again. We can see that the positive root spaces will correspond to $e_{i,j}$ for $i < j$. So if we choose e_α s for the positive root α to be the $e_{i,j}$ in $\mathfrak{g}_\alpha$, we will have $i < j$ and $e_{-\alpha}$ will be $(-1)^{[i]} e_{j,i}$. Then we will have:*

$$\begin{aligned} e_\alpha^* &= e_{i,j}^* = (-1)^{[i]} e_{j,i} = e_{-\alpha} \\ e_{-\alpha}^* &= (-1)^{[i]} e_{j,i}^* = (-1)^{[i]} (-1)^{[j]} e_{i,j} = (-1)^{|\alpha|} e_\alpha \end{aligned}$$

and the above formula for Ω will agree with the Casimir element found earlier.

⁶This will hold for all classical Lie superalgebras of the form: $A(m, n)$ for $(m, n) \neq (1, 1)$, $B(m, n)$, $C(n)$, $D(m, n)$, $D(2, 1; \alpha)$, $F(4)$, and $G(3)$. In fact the dimension of $\mathfrak{g}_\alpha$ is one provided $\mathfrak{g}$ is a classical Lie superalgebra different from $A(1, 1)$, $P(2)$, $P(3)$, and $Q(n)$. See [4] for more on Lie superalgebras.

3.4. Statement of the Theorem: We are now ready to state our main theorem. Its proof will be presented in the next three sections.

Theorem 2. *Let $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ satisfy:*

$$r_0^{12} + r_0^{21} = \Omega_0 \quad (1)$$

$$(\tau(\alpha) \otimes 1)(r_0) + (1 \otimes \alpha)(r_0) = 0 \text{ for all } \alpha \in \Gamma_1 \quad (2)$$

Then the element r of $\mathfrak{g} \otimes \mathfrak{g}$ defined by:

$$r = r_0 + \sum_{\alpha > 0} e_{-\alpha} \otimes e_{\alpha} + \sum_{\alpha, \beta > 0, \alpha < \beta} (e_{-\alpha} \otimes e_{\beta} - (-1)^{|\alpha|} e_{\beta} \otimes e_{-\alpha}) \quad (*)$$

is a solution to the system:

$$r^{12} + r^{21} = \Omega \quad (3)$$

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0 \quad (4)$$

Remark. Comparing this theorem with the corresponding theorem for Lie algebras, we notice that there is a sign introduced in the equation defining the r -matrix associated to a given admissible triple. However since Lie algebras are Lie superalgebras with only even roots, Equation $(*)$ reduces to the corresponding equation in Theorem 1 when the Lie superalgebra $\mathfrak{g}$ in question is actually a Lie algebra.

4. TECHNICAL LEMMAS

Let $\mathfrak{g}$ be a simple Lie superalgebra with a non-degenerate Killing form ⁷. Assume that we fix a homogeneous basis $\{I_{\alpha}\}$ for $\mathfrak{g}$ and denote by $\{I_{\alpha}^*\}$ the dual basis of $\mathfrak{g}$ with respect to the non-degenerate (Killing) form. We will first prove:

Lemma 1. *Let $r \in \mathfrak{g} \otimes \mathfrak{g}$ be such that*

$$r = (f \otimes 1)\Omega$$

Then the system of equations:

$$r^{12} + r^{21} = \Omega \quad (3)$$

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0 \quad (4)$$

is equivalent to the system:

$$f + f^* = 1 \quad (5)$$

$$(f - 1)[f(x), f(y)] = f([(f - 1)(x), (f - 1)(y)]) \quad (6)$$

where f^ stands for the adjoint of f with respect to the standard form $(\ , \)$.*

Remark. This is exactly the same technical lemma used in the proof of the main classification theorem in the Lie algebra case as presented in [3]. Our proof here will be a generalization of the proof provided there. We will use general properties like the invariance, non-degeneracy and the consistency of the Killing form. The main difference will be that in our case, we may not be able to pick an orthonormal basis for our Lie superalgebra $\mathfrak{g}$. However it turns out that a pair of dual bases will be sufficient for our purposes.

⁷Our results can mostly be extended to the class of classical Lie superalgebras. If $\mathfrak{g}$ is classical, there is a non-degenerate invariant form on $\mathfrak{g}$. In the following, most of the statements involving the Killing form may be asserted more generally for such an invariant form.

Proof: PART 1: [This part is exactly the same as in the Lie algebra case.] We have:

$$r^{12} + r^{21} = (f \otimes 1)\Omega + (1 \otimes f)\Omega = (f \otimes 1)\Omega + (f^* \otimes 1)\Omega = ((f + f^*) \otimes 1)\Omega.$$

Then we have:

$$\begin{aligned} \Omega &= r^{12} + r^{21} \\ &\Leftrightarrow \\ \Omega &= (1 \otimes 1)\Omega = ((f + f^*) \otimes 1)\Omega \\ &\Leftrightarrow \\ 1 &= (f + f^*) \end{aligned}$$

This proves the equivalence of the statements:

$$\Omega = r^{12} + r^{21} \quad \text{and} \quad 1 = (f + f^*)$$

PART 2: Next we show that the CYBE for r (that is, Equation 4), translates to a nice expression in terms of the associated function f . This part of the proof requires some modifications to the proof of the Lie algebra case.

We have:

$$r = (f \otimes 1)\Omega = \sum_{\alpha} (-1)^{|I_{\alpha}|} f(I_{\alpha}) \otimes I_{\alpha}^*$$

Let us write the three terms of the CYBE:

$$\begin{aligned} [r^{12}, r^{13}] &= \sum_{\alpha, \beta} (-1)^{|I_{\alpha}|+|I_{\beta}|} (-1)^{|I_{\alpha}||I_{\beta}|} [f(I_{\alpha}), f(I_{\beta})] \otimes I_{\alpha}^* \otimes I_{\beta}^* \\ [r^{12}, r^{23}] &= \sum_{\alpha, \beta} (-1)^{|I_{\alpha}|+|I_{\beta}|} f(I_{\alpha}) \otimes [I_{\alpha}^*, f(I_{\beta})] \otimes I_{\beta}^* \\ [r^{13}, r^{23}] &= \sum_{\alpha, \beta} (-1)^{|I_{\alpha}|+|I_{\beta}|} (-1)^{|I_{\alpha}||I_{\beta}|} f(I_{\alpha}) \otimes f(I_{\beta}) \otimes [I_{\alpha}^*, I_{\beta}^*] \end{aligned}$$

Here we use:

$$\begin{aligned} [a \otimes b \otimes 1, c \otimes 1 \otimes d] &= (-1)^{|b||c|} [a, c] \otimes b \otimes d \\ [a \otimes b \otimes 1, 1 \otimes c \otimes d] &= a \otimes [b, c] \otimes d \\ [a \otimes 1 \otimes b, 1 \otimes c \otimes d] &= (-1)^{|b||c|} a \otimes c \otimes [b, d] \end{aligned}$$

and:

$$|I_{\alpha}| = |I_{\alpha}^*| \quad \text{and} \quad |f(I_{\alpha})| = |I_{\alpha}|$$

(We assume f is even.) We rewrite the last sum so that it ends with $\otimes I_{\beta}^*$:

$$\begin{aligned} &\sum_{\alpha, \beta} (-1)^{|I_{\alpha}|+|I_{\beta}|} (-1)^{|I_{\alpha}||I_{\beta}|} f(I_{\alpha}) \otimes f(I_{\beta}) \otimes [I_{\alpha}^*, I_{\beta}^*] \\ &= - \sum_{\alpha, \beta} (-1)^{|I_{\alpha}|+|I_{\beta}|} f(I_{\alpha}) \otimes f([I_{\alpha}^*, I_{\beta}]) \otimes I_{\beta}^* \end{aligned}$$

where we use the invariance of the form, and the supersymmetry of the bracket.

Therefore we can rewrite the CYBE as:

$$\sum_{\alpha, \beta} (-1)^{|I_{\beta}|} \left(\begin{aligned} &(-1)^{|I_{\alpha}|} (-1)^{|I_{\alpha}||I_{\beta}|} [f(I_{\alpha}), f(I_{\beta})] \otimes I_{\alpha}^* \\ &+ (-1)^{|I_{\alpha}|} f(I_{\alpha}) \otimes [I_{\alpha}^*, f(I_{\beta})] \\ &- (-1)^{|I_{\alpha}|} f(I_{\alpha}) \otimes f([I_{\alpha}^*, I_{\beta}]) \end{aligned} \right) \otimes I_{\beta}^* = 0$$

Since the $\{I_\beta^*\}$ form a basis for $\mathfrak{g}$, this last equation implies that, for any choice of β , we have:

$$\begin{pmatrix} \sum_{\alpha} (-1)^{|I_\alpha|} (-1)^{|I_\alpha||I_\beta|} [f(I_\alpha), f(I_\beta)] \otimes I_\alpha^* \\ + \sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes [I_\alpha^*, f(I_\beta)] \\ - \sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes f([I_\alpha^*, I_\beta]) \end{pmatrix} = 0$$

We want to rewrite the second and the third sums so that they end with $\otimes I_\alpha^*$. After some calculation, the second term becomes:

$$\sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes [I_\alpha^*, f(I_\beta)] = \boxed{- \sum_{\alpha} (-1)^{|I_\alpha|} (-1)^{|I_\alpha||I_\beta|} f([I_\alpha, f(I_\beta)]) \otimes I_\alpha^*}$$

The third sum splits into two different sums when we use Equation 3:

$$\begin{aligned} & - \sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes f([I_\alpha^*, I_\beta]) \\ & = - \sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes [I_\alpha^*, I_\beta] + \sum_{\alpha} (-1)^{|I_\alpha|} f(f(I_\alpha)) \otimes [I_\alpha^*, I_\beta] \end{aligned}$$

We calculate these terms separately:

$$- \sum_{\alpha} (-1)^{|I_\alpha|} f(I_\alpha) \otimes [I_\alpha^*, I_\beta] = \boxed{\sum_{\alpha} (-1)^{|I_\alpha|} (-1)^{|I_\alpha||I_\beta|} f([I_\alpha, I_\beta]) \otimes I_\alpha^*}$$

and similar but more difficult calculations yield:

$$\sum_{\alpha} (-1)^{|I_\alpha|} f(f(I_\alpha)) \otimes [I_\alpha^*, I_\beta] = \boxed{- \sum_{\alpha} (-1)^{|I_\alpha|} (-1)^{|I_\alpha||I_\beta|} f([f(I_\alpha), I_\beta]) \otimes I_\alpha^*}$$

Hence we get:

$$\sum_{\alpha} (-1)^{|I_\alpha|} (-1)^{|I_\alpha||I_\beta|} \begin{pmatrix} [f(I_\alpha), f(I_\beta)] & - & f([I_\alpha, f(I_\beta)]) \\ +f([I_\alpha, I_\beta]) & - & f([f(I_\alpha), I_\beta]) \end{pmatrix} \otimes I_\alpha^* = 0$$

Again using the fact that the $\{I_\alpha^*\}$ form a basis for $\mathfrak{g}$, we obtain, for all α, β :

$$\begin{aligned} & [f(I_\alpha), f(I_\beta)] - f([I_\alpha, f(I_\beta)]) + f([I_\alpha, I_\beta]) - f([f(I_\alpha), I_\beta]) = 0 \\ \Rightarrow & [f(I_\alpha), f(I_\beta)] = f([I_\alpha, f(I_\beta)]) - f([I_\alpha, I_\beta]) + f([f(I_\alpha), I_\beta]) \\ \Rightarrow & (f - 1)[f(I_\alpha), f(I_\beta)] = f([(f - 1)(I_\alpha), (f - 1)(I_\beta)]), \end{aligned}$$

which is equivalent to

$$\boxed{(f - 1)[f(x), f(y)] = f([(f - 1)(x), (f - 1)(y)]) \text{ for all } x, y \in \mathfrak{g}} \quad (6)$$

This proves one direction of the lemma. To see the other direction, we need only trace the steps above backwards. Hence one can easily see that a function f satisfying Equations 5 and 6 will correspond to an r -matrix $r \in \mathfrak{g} \otimes \mathfrak{g}$ that satisfies Equations 3 and 4. This completes the proof. ■

Writing $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ as $r_0 = (f_0 \otimes 1)\Omega_0$ for a linear map $f_0 : \mathfrak{h} \rightarrow \mathfrak{h}$, we have:

Lemma 2. *The system:*

$$r_0^{12} + r_0^{21} = \Omega_0 \quad (1)$$

$$(\tau(\alpha) \otimes 1)(r_0) + (1 \otimes \alpha)(r_0) = 0 \text{ for all } \alpha \in \Gamma_1 \quad (2)$$

is equivalent to the system:

$$f_0 + f_0^* = 1 \quad (7)$$

$$f_0(h_\alpha) = (f_0 - 1)(h_{\tau(\alpha)}) \text{ for all } \alpha \in \Gamma_1 \quad (8)$$

Proof: We will prove a stronger result. Namely, we will prove that, for any $1 \leq s, t \leq r$, the equations:

$$r_0^{12} + r_0^{21} = \Omega_0$$

$$(\alpha_t \otimes 1)(r_0) + (1 \otimes \alpha_s)(r_0) = 0$$

are equivalent to the equations:

$$f_0 + f_0^* = 1$$

$$f_0(h_{\alpha_s}) = (f_0 - 1)(h_{\alpha_t})$$

It is easy to see the equivalence of the first equations:

$$\begin{aligned} r_0^{12} + r_0^{21} &= (f_0 \otimes 1 + 1 \otimes f_0)\Omega_0 \\ &= ((f_0 + f_0^*) \otimes 1)\Omega_0 \\ &= \Omega_0 \text{ if and only if } f_0 + f_0^* = 1 \end{aligned}$$

Next we look at $(\alpha_t \otimes 1)r_0 + (1 \otimes \alpha_s)r_0$. This is equal to:

$$\begin{aligned} &(\alpha_t \otimes 1) \left(\sum_i f_0(h_{\alpha_i}) \otimes h_{\alpha_i}^* \right) + (1 \otimes \alpha_s) \left(\sum_i f_0(h_{\alpha_i}) \otimes h_{\alpha_i}^* \right) \\ &= \sum_i \alpha_t(f_0(h_{\alpha_i})) \cdot h_{\alpha_i}^* + \sum_i \alpha_s(f_0^*(h_{\alpha_i}^*)) \cdot h_{\alpha_i} \\ &= \sum_i \alpha_t \left(\sum_k (f_0(h_{\alpha_i}), h_{\alpha_k}) h_{\alpha_k}^* \right) \cdot h_{\alpha_i}^* + \sum_i \alpha_s \left(\sum_k (f_0^*(h_{\alpha_i}^*), h_{\alpha_k}) h_{\alpha_k}^* \right) \cdot h_{\alpha_i} \\ &= \sum_{i,k} (f_0(h_{\alpha_i}), h_{\alpha_k}) \alpha_t(h_{\alpha_k}^*) \cdot h_{\alpha_i}^* + \sum_{i,k} (f_0^*(h_{\alpha_i}^*), h_{\alpha_k}) \alpha_s(h_{\alpha_k}^*) \cdot h_{\alpha_i} \end{aligned}$$

We have:

$$\alpha_s(h_{\alpha_k}^*) = (h_{\alpha_s}, h_{\alpha_k}^*) = \delta_{s,k} \quad \text{and} \quad \alpha_t(h_{\alpha_k}^*) = (h_{\alpha_t}, h_{\alpha_k}^*) = \delta_{t,k}$$

Therefore the above expression becomes:

$$\begin{aligned}
& \sum_i (f_0(h_{\alpha_i}), h_{\alpha_t}) h_{\alpha_i}^* + \sum_i (f_0^*(h_{\alpha_i}^*), h_{\alpha_s}) h_{\alpha_i} \\
&= \sum_i (h_{\alpha_i}, f_0^*(h_{\alpha_t})) h_{\alpha_i}^* + \sum_i (h_{\alpha_i}^*, f_0(h_{\alpha_s})) h_{\alpha_i} \\
&= f_0^*(h_{\alpha_t}) + f_0(h_{\alpha_s}) \\
&= (1 - f_0)(h_{\alpha_t}) + f_0(h_{\alpha_s})
\end{aligned}$$

Summarizing, we have shown that:

$$(\alpha_t \otimes 1)r_0 + (1 \otimes \alpha_s)r_0 = (1 - f_0)(h_{\alpha_t}) + f_0(h_{\alpha_s})$$

Hence one is equal to zero if and only if the other is. This completes the proof of the lemma. ■

We also need:

Lemma 3. *The system of equations*

$$r_0^{12} + r_0^{21} = \Omega_0 \tag{1}$$

$$(\tau(\alpha) \otimes 1)(r_0) + (1 \otimes \alpha)(r_0) = 0 \text{ for all } \alpha \in \Gamma_1 \tag{2}$$

is consistent.

Remark. *The arguments used to prove this lemma are the same as for the Lie algebra case (see [2] for details), and hence will not be included here.*

These lemmas allow us to translate the conditions (Equations 1 and 2) on the continuous parameter used in the main theorem into conditions on a linear map $f_0 : \mathfrak{h} \rightarrow \mathfrak{h}$. Also recall that Lemma 1 translates the CYBE and the unitarity condition (i.e. Equations 3 and 4) into conditions on the associated linear map $f : \mathfrak{g} \rightarrow \mathfrak{g}$, namely Equations 5 and 6.

So from now on, we will be using linear maps f, f_0 and so on, interchangeably with their 2-tensor equivalents r, r_0 etc. Then, we can reformulate our problem in the following manner: Given an admissible triple $(\Gamma_1, \Gamma_2, \tau)$ with a linear map $f_0 : \mathfrak{h} \rightarrow \mathfrak{h}$ satisfying Equations 7 and 8, construct a linear map $f : \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying Equations 5 and 6.

5. THE CAYLEY TRANSFORM

We will now consider a variation on the theme of Cayley transforms. For the time being assume that we have a linear function $f : \mathfrak{g} \rightarrow \mathfrak{g}$ with $(f - 1)$ invertible. Then **the Cayley transform of f** is $\Theta = \frac{f}{f-1}$. Then the adjoint of this function will be $\Theta^* = \frac{f^*}{(f-1)^*} = \frac{1-f}{-f}$ if our f satisfies Equation 5:

$$f + f^* = 1 \tag{5}$$

Then one can see that $\Theta\Theta^* = 1$, and so Θ preserves the invariant form. If we also assume that f satisfies Equation 6:

$$(f - 1)([f(x), f(y)]) = f([(f - 1)(x), (f - 1)(y)]) \tag{6}$$

we will get:

$$[\Theta(x), \Theta(y)] = \Theta([x, y])$$

which implies that Θ is a Lie superalgebra automorphism.

However for the Lie superalgebras we care about, i.e. for simple Lie superalgebras, the above will not work. To see this, assume that f is a linear map satisfying Equations 5 and 6, $f - 1$ is invertible, and Θ is defined as above. Then look at $\Theta - 1$. This is given by $\frac{f}{f-1} - 1 = \frac{1}{f-1}$, i.e. it is the inverse of $f - 1$. So we must have that $\det(\Theta - 1) \neq 0$. But this is not allowed for a simple Lie superalgebra:

Lemma 4. *If Θ is an automorphism of a finite dimensional (classical) simple Lie superalgebra, then $\det(\Theta - 1) = 0$.*

Proof: We will need the following result from [2] (this is Theorem 9.2 there):

If φ is an automorphism of a semisimple Lie algebra L , then there exists a nonzero element $x \in L$ such that $\varphi(x) = x$.

Now the automorphism Θ of a simple Lie superalgebra $\mathfrak{g}$ restricts to a (Lie algebra) automorphism θ on $\mathfrak{g}_0$, the even part of $\mathfrak{g}$. $\mathfrak{g}_0$ is reductive with nontrivial $\mathfrak{g}_0' = [\mathfrak{g}_0, \mathfrak{g}_0]$ ⁸. $\mathfrak{g}_0'$ is semisimple and θ restricts to an automorphism φ on $\mathfrak{g}_0'$. But then the above result gives us some nonzero $x \in \mathfrak{g}_0'$ with $\varphi(x) = x$. Going back to our Θ we see that $\Theta(x) = x$ and hence $x \in \text{Ker}(\Theta - 1)$. Thus we must have that $\det(\Theta - 1) = 0$. This proves the lemma. ■

Thus Equations 5 and 6 will imply that $f - 1$ is not invertible, as the invertibility assumption leads to a contradiction with Lemma 4⁹. Therefore, we cannot define the Cayley transform as above for the functions we are interested in.

However it turns out that we can modify our definition and still get a lot of what we want: First note that for any linear operator f , $\text{Ker}(f) \subset \text{Im}(f - 1)$ and $\text{Ker}(f - 1) \subset \text{Im}(f)$. Then we define the **Cayley transform of f** to be the function $\Theta : \text{Im}(f - 1)/\text{Ker}(f) \rightarrow \text{Im}(f)/\text{Ker}(f - 1)$ that maps $\overline{(f - 1)(x)}$ to $\overline{f(x)}$. [It is easy to check that this is well-defined.] This version of the Cayley transform will be sufficient for our purposes. We have:

Lemma 5. *Let $f : \mathfrak{g} \rightarrow \mathfrak{g}$ be a linear map satisfying:*

$$f + f^* = 1 \tag{5}$$

Then $\text{Ker}(f) = \text{Im}(f - 1)^\perp$, $\text{Ker}(f - 1) = \text{Im}(f)^\perp$, and the map Θ preserves the invariant form. Furthermore, f satisfies:

$$(f - 1)[f(x), f(y)] = f([(f - 1)(x), (f - 1)(y)]), \tag{6}$$

if and only if $\text{Im}(f)$ and $\text{Im}(f - 1)$ are Lie subsuperalgebras of $\mathfrak{g}$, and Θ is a Lie superalgebra isomorphism.

Remark. *This lemma will be valid for any simple Lie superalgebra $\mathfrak{g}$ with a non-degenerate Killing form. In this case its proof will be exactly the same as the proof of the analogous result in the Lie algebra case. See [2].*

⁸The even part of a classical simple Lie superalgebra will be reductive and will decompose into a direct sum of its derived algebra (which is nonempty) and some abelian subalgebra.

⁹Using Equation 5 once again, we see that f is not invertible, either. Thus any solution r to the system of Equations 5 and 6 will be degenerate.

6. THE CONSTRUCTION - END OF THE PROOF OF THE THEOREM

For a given admissible triple $(\Gamma_1, \Gamma_2, \tau)$, and a linear map $f_0 : \mathfrak{h} \rightarrow \mathfrak{h}$ satisfying Equations 7 and 8, we want to construct a function $f : \mathfrak{g} \rightarrow \mathfrak{g}$ that will satisfy Equations 5 and 6. Here is how we proceed:

For the admissible triple $(\Gamma_1, \Gamma_2, \tau)$, we define $\bar{\Gamma}_i$ and $\bar{\tau}$ as before. Also we define the following Lie subsuperalgebras of $\mathfrak{g}$:

$$\mathfrak{h}_i = \bigoplus_{\alpha \in \Gamma_i} \mathbb{C} h_\alpha \quad \mathfrak{g}_i = \mathfrak{h}_i \oplus \sum_{\alpha \in \bar{\Gamma}_i} (\mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha})$$

and:

$$\begin{aligned} \mathfrak{n}_i^+ &= \sum_{\alpha \in \Delta^+ / \bar{\Gamma}_i} \mathfrak{g}_\alpha & \mathfrak{p}_i^+ &= \mathfrak{g}_i + \mathfrak{n}_i^+ \\ \mathfrak{n}_i^- &= \sum_{\alpha \in \Delta^+ / \bar{\Gamma}_i} \mathfrak{g}_{-\alpha} & \mathfrak{p}_i^- &= \mathfrak{g}_i + \mathfrak{n}_i^- \end{aligned}$$

We can see that the $\mathfrak{n}_i^{+/-}$ are ideals in $\mathfrak{p}_i^{+/-}$.

Next if $f_0 : \mathfrak{h} \rightarrow \mathfrak{h}$ satisfies:

$$f_0(h_\alpha) = (f_0 - 1)(h_{\tau(\alpha)}) \text{ for all } \alpha \in \Gamma_1 \quad (8)$$

we get:

$$h_\alpha = (f_0 - 1)(h_{\tau(\alpha)} - h_\alpha) \quad h_{\tau(\alpha)} = f_0(h_{\tau(\alpha)} - h_\alpha)$$

for all $\alpha \in \Gamma_1$. This implies that $h_\alpha \in \text{Im}(f_0 - 1)$ and $h_{\tau(\alpha)} \in \text{Im}(f_0)$. Therefore we have that: $\mathfrak{h}_1 \subset \text{Im}(f_0 - 1)$, and $\mathfrak{h}_2 \subset \text{Im}(f_0)$.

Now fix a Weyl-Chevalley basis $\{X_{\alpha_i}, Y_{\alpha_i}, H_{\alpha_i} | \alpha_i \in \Gamma\}$. It is known that such a set of generators exists and satisfies the usual Serre-type relations, (see [5] for details). We define a map φ by

$$\varphi(X_\alpha) = X_{\tau(\alpha)} \quad \varphi(Y_\alpha) = Y_{\tau(\alpha)} \quad \varphi(H_\alpha) = H_{\tau(\alpha)}$$

for all $\alpha \in \Gamma_1$. Then this can be extended to an isomorphism $\varphi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ because the relations between $X_\alpha, Y_\alpha, H_\alpha$ for $\alpha \in \Gamma_1$ will be the same as the relations between $X_{\tau(\alpha)}, Y_{\tau(\alpha)}, H_{\tau(\alpha)}$ for $\alpha \in \Gamma_1$ [Here we are using the fact that τ is an isometry preserving grading]. Note that φ^{-1} is a map from $\mathfrak{g}_2$ onto $\mathfrak{g}_1$. Since τ is an isometry, $(\varphi(x), y)_{\mathfrak{g}_2} = (x, \varphi^{-1}(y))_{\mathfrak{g}_1}$ for all $x \in \mathfrak{g}_1, y \in \mathfrak{g}_2$. But φ^* should map $\mathfrak{g}_2$ into $\mathfrak{g}_1$ and satisfy exactly the same conditions; hence $\varphi^* = \varphi^{-1}$.

Now in each root space $\mathfrak{g}_\alpha$, we can choose an element e_α such that $(e_\alpha, e_{-\alpha}) = 1$ for any α and $\varphi(e_\alpha) = e_{\bar{\tau}(\alpha)}$ for all $\alpha \in \bar{\Gamma}_1$. The fact that there are no cycles for τ will ensure that such a choice is possible.

Next define a linear map as follows:

$$\psi(x) = \begin{cases} \varphi(x) & \text{if } x \in \mathfrak{g}_1 \\ 0 & \text{if } x \in \mathfrak{n}_1^+ \end{cases}$$

This restricts to a map on $\mathfrak{n}_+ = \bigoplus_{\alpha > 0} \mathfrak{g}_\alpha$, since $\mathfrak{n}_+ = (\mathfrak{g}_1 \cap \mathfrak{n}_+) \oplus \mathfrak{n}_1^+$. The proof of the following lemma is exactly the same as in the Lie algebra case (see [2]):

Lemma 6. *det($\psi - 1$) is nonzero if and only if τ satisfies the second condition in the definition of an admissible triple.*

Recall that we started with an admissible triple. The above lemma then says that $\psi - 1$ is invertible. Therefore we can define a function on $\mathfrak{n}_+$ by:

$$f_+ = \frac{\psi}{\psi - 1} = -(\psi + \psi^2 + \cdots)$$

Clearly the sum on the right hand side is finite as ψ is nilpotent. Notice that ψ^* and so f_+^* are maps on $\mathfrak{n}_- = \bigoplus_{\alpha < 0} \mathfrak{g}_\alpha$, since the Killing form induces a non-degenerate pairing of $\mathfrak{n}_+$ with $\mathfrak{n}_-$.

Now define a linear map on $\mathfrak{n}_-$ by:

$$f_- = 1 - f_+^* = 1 + \psi^* + \psi^{*2} + \cdots$$

Then define f to be the function whose restriction to $\mathfrak{h}$, $\mathfrak{n}_+$, $\mathfrak{n}_-$ is f_0 , f_+ , f_- , respectively. We have:

$$\begin{aligned} f + f^* &= (f_0 + f_+ + f_-) + (f_0 + f_+ + f_-)^* \\ &= (f_0 + f_0^*) + (f_+ + f_-^*) + (f_+^* + f_-) \\ &= 1_{\mathfrak{h}} + 1_{\mathfrak{n}_+} + 1_{\mathfrak{n}_-} \\ &= 1_{\mathfrak{g}} \end{aligned}$$

To see that f satisfies Equation 6, we will use Lemma 5. Recall that for a linear map $f : \mathfrak{g} \rightarrow \mathfrak{g}$, we defined the Cayley transform to be the function

$$\Theta : Im(f - 1)/Ker(f) \rightarrow Im(f)/Ker(f - 1)$$

that maps $\overline{(f - 1)(x)}$ to $\overline{f(x)}$. Then we have seen before that if f satisfies Equation 5 i.e. $f + f^* = 1$, then $Ker(f) = Im(f - 1)^\perp$, $Ker(f - 1) = Im(f)^\perp$, and $\Theta\Theta^* = 1$. Furthermore, f satisfies Equation 6 if and only if $Im(f)$ and $Im(f - 1)$ are Lie subsuperalgebras of $\mathfrak{g}$, and Θ is a Lie superalgebra isomorphism.

Thus our problem now reduces to showing that $C_1 = Im(f - 1)$ and $C_2 = Im(f)$ are Lie subsuperalgebras of $\mathfrak{g}$, and that the Cayley transform Θ of f is a Lie superalgebra isomorphism.

We have

$$\begin{aligned} C_1 &= Im(f - 1) = Im(f_0 - 1) \cup Im(f_+ - 1) \cup Im(f_- - 1) \\ C_2 &= Im(f) = Im(f_0) \cup Im(f_+) \cup Im(f_-) \end{aligned}$$

We have seen that $Im(f_0 - 1) \supset \mathfrak{h}_1$ and $Im(f_0) \supset \mathfrak{h}_2$. We will therefore define V_1 , V_2 as (vector) subspaces of $\mathfrak{h}$ such that $Im(f_0 - 1) = \mathfrak{h}_1 \oplus V_1$ and $Im(f_0) = \mathfrak{h}_2 \oplus V_2$.

In the Lie algebra case, the Killing form restricts to a positive definite non-degenerate form on (the real subspace generated by $\{H_\alpha | \alpha \in \Gamma\}$ of) $\mathfrak{h}$. So we can define the orthogonal complements of $\mathfrak{h}_1$ and $\mathfrak{h}_2$ with respect to this form; call these $\mathfrak{h}_1^c$ and $\mathfrak{h}_2^c$; then we have: $\mathfrak{h} = \mathfrak{h}_1 \oplus \mathfrak{h}_1^c = \mathfrak{h}_2 \oplus \mathfrak{h}_2^c$. Then for a fixed f_0 , the two subspaces V_1 and V_2 are uniquely determined if we add the condition that $V_i \subset \mathfrak{h}_i^c$.

In the super case, this is no longer possible; the Cartan subalgebra $\mathfrak{h}$ may have isotropic elements and subspaces of $\mathfrak{h}$ may intersect their orthogonal complements nontrivially. However in our case we still can define $\mathfrak{h}_i^c$ as follows:

$$\mathfrak{h}_i^c = \bigoplus_{\alpha \in \Gamma \setminus \Gamma_i} \mathbb{C}h_\alpha$$

Thus we still can write $\mathfrak{h} = \mathfrak{h}_i \oplus \mathfrak{h}_i^c$, and still can demand that $V_i \subset \mathfrak{h}_i^c$. This choice of the V_i is then again well-defined, but clearly depends on our choice for Γ .

Next we compute:

$$\begin{aligned} \text{Im}(f_+ - 1) &= \text{Im}\left(\frac{1}{\psi - 1}\right) = \mathfrak{n}_+ \\ \text{Im}(f_- - 1) &= \text{Im}\left(\frac{\psi^*}{1 - \psi^*}\right) = \text{Im}(\psi^*) = \mathfrak{g}_1 \cap \mathfrak{n}_- \\ \text{Im}(f_+) &= \text{Im}\left(\frac{\psi}{1 - \psi}\right) = \text{Im}(\psi) = \mathfrak{g}_2 \cap \mathfrak{n}_+ \\ \text{Im}(f_-) &= \text{Im}\left(\frac{1}{1 - \psi^*}\right) = \mathfrak{n}_- \end{aligned}$$

where we use the fact that $\psi - 1$ is invertible. The above then yields:

$$C_1 = \mathfrak{p}_1^+ \oplus V_1 \quad C_2 = \mathfrak{p}_2^- \oplus V_2$$

It is now easy to check that C_1 and C_2 are both closed under the bracket and hence are Lie subsuperalgebras of $\mathfrak{g}$.

Finally we need to see that the Cayley transform Θ is a Lie superalgebra isomorphism. Now we note that by the last lemma above, $C_i \supset C_i^\perp$. So we have:

$$C_1^\perp = (\mathfrak{p}_1^+ \oplus V_1)^\perp = \mathfrak{n}_1^+ \oplus (\mathfrak{h}_1 \oplus V_1)^\perp = \mathfrak{n}_1^+ \oplus (\mathfrak{h}_1^\perp \cap V_1^\perp) \subset \mathfrak{p}_1^+ \oplus V_1$$

and similarly:

$$C_2^\perp = (\mathfrak{p}_2^- \oplus V_2)^\perp = \mathfrak{n}_2^- \oplus (\mathfrak{h}_2 \oplus V_2)^\perp = \mathfrak{n}_2^- \oplus (\mathfrak{h}_2^\perp \cap V_2^\perp) \subset \mathfrak{p}_2^- \oplus V_2$$

Hence we have:

$$\mathfrak{h}_i^\perp \cap V_i^\perp \subset \mathfrak{h}_i \oplus V_i$$

which gives us:

$$C_1/C_1^\perp = \frac{\mathfrak{p}_1^+ \oplus V_1}{(\mathfrak{p}_1^+ \oplus V_1)^\perp} = \left(\bigoplus_{\alpha \in \Gamma_1} \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \right) \oplus \frac{\mathfrak{h}_1 \oplus V_1}{\mathfrak{h}_1^\perp \cap V_1^\perp}$$

and similarly:

$$C_2/C_2^\perp = \frac{\mathfrak{p}_2^- \oplus V_2}{(\mathfrak{p}_2^- \oplus V_2)^\perp} = \left(\bigoplus_{\alpha \in \Gamma_2} \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \right) \oplus \frac{\mathfrak{h}_2 \oplus V_2}{\mathfrak{h}_2^\perp \cap V_2^\perp}$$

We have already seen that the C_i are Lie subsuperalgebras. Since $C_i^\perp$ is an ideal, we have a Lie superalgebra structure on $C_i/C_i^\perp$. But $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] = \mathbb{C}H_\alpha$, therefore we must have a complete copy of $\mathfrak{h}_i$ and so a copy of $\mathfrak{g}_i$ in $C_i/C_i^\perp$. This implies that $\mathfrak{h}_i^\perp \cap V_i^\perp$ lies in V_i , and we have:

$$C_i/C_i^\perp = \mathfrak{g}_i \oplus \frac{V_i}{\mathfrak{h}_i^\perp \cap V_i^\perp}$$

So we need to show that:

$$\Theta : \mathfrak{g}_1 \oplus \frac{V_1}{\mathfrak{h}_1^\perp \cap V_1^\perp} \rightarrow \mathfrak{g}_2 \oplus \frac{V_2}{\mathfrak{h}_2^\perp \cap V_2^\perp}$$

is a Lie superalgebra isomorphism.

We first note that $\Theta(x) = \varphi(x)$ for all $x \in \mathfrak{g}_1$. Indeed if $\alpha \in \Gamma_1$, we have:

$$X_\alpha = (f_+ - 1)(X_{\tau(\alpha)} - X_\alpha)$$

and so is mapped via Θ to:

$$f_+(X_{\tau(\alpha)} - X_\alpha) = X_{\tau(\alpha)}$$

And similarly:

$$Y_\alpha = (f_- - 1)(Y_{\tau(\alpha)} - Y_\alpha)$$

is mapped via Θ to:

$$f_-(Y_{\tau(\alpha)} - Y_\alpha) = Y_{\tau(\alpha)}$$

Also it is easy to see that since $H_\alpha = (f_0 - 1)(H_{\tau(\alpha)} - H_\alpha)$ for each $\alpha \in \Gamma_1$, Θ sends H_α to $f_0(H_{\tau(\alpha)} - H_\alpha) = H_{\tau(\alpha)}$. Hence the restriction of Θ to $\mathfrak{g}_1$ is exactly the Lie superalgebra isomorphism φ .

Next we look at what Θ does on the Cartan part of the $C_i/C_i^\perp$. We have that:

$$C_i/C_i^\perp = \left(\bigoplus_{\alpha \in \Gamma_i} \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \right) \oplus \frac{\mathfrak{h}_i \oplus V_i}{\mathfrak{h}_i^\perp \cap V_i^\perp}$$

i.e. we can rewrite $C_i/C_i^\perp$ as the direct sum of a Cartan part and a non-Cartan part. Then the previous arguments show that Θ maps the non-Cartan part of $C_1/C_1^\perp$ into the non-Cartan part of $C_2/C_2^\perp$ like φ does. And then since Θ preserves the invariant form, it maps the Cartan part of $C_1/C_1^\perp$ to the Cartan part of $C_2/C_2^\perp$:

$$\Theta \left((\text{non-Cartan of } C_1/C_1^\perp)^\perp \right) = (\text{non-Cartan of } C_2/C_2^\perp)^\perp$$

In other words:

$$\Theta \left(\frac{\mathfrak{h}_1 \oplus V_1}{\mathfrak{h}_1^\perp \cap V_1^\perp} \right) = \left(\frac{\mathfrak{h}_2 \oplus V_2}{\mathfrak{h}_2^\perp \cap V_2^\perp} \right)$$

Since $\frac{\mathfrak{h}_i \oplus V_i}{\mathfrak{h}_i^\perp \cap V_i^\perp}$ are abelian, Θ restricts to an isomorphism there as well. Therefore Θ is an isomorphism. This then will imply that the associated linear map f satisfies Equations 5 and 6 and so corresponds to an r -matrix satisfying Equations 3 and 4.

At this point one needs to check if the function f constructed in this way will yield the tensor r of Equation (*). This is reasonably straightforward. Hence we have proved our theorem. ■

7. EXAMPLES: r -MATRICES ON $sl(2,1)$

Recall that two Dynkin diagrams of a given Lie superalgebra are not necessarily isomorphic, but one can be obtained from the other via a chain of odd reflections. Therefore while listing all possible admissible triples for a given Lie superalgebra, we need to take into consideration all possible Dynkin diagrams. This raises a new question as to how r -matrices obtained from two nonisomorphic Dynkin diagrams are related, if at all. We will see that at least in the case of $sl(2,1)$, if r is the standard r -matrix associated to a fixed Dynkin diagram D , and D' is the Dynkin diagram obtained from D by the odd reflection σ_α , then r' , the standard r -matrix associated to D' , will be the image of r under the same reflection σ_α .

7.1. Dynkin Diagrams of $sl(2,1)$. The roots of $sl(2,1)$ are

$$\Delta_{\overline{0}} = \{\epsilon_1 - \epsilon_2, \epsilon_2 - \epsilon_1\} \quad \Delta_{\overline{1}} = \{\epsilon_1 - \lambda_1, \epsilon_2 - \lambda_1, \lambda_1 - \epsilon_1, \lambda_1 - \epsilon_2\}$$

where ϵ_i is the (restriction to the Cartan subalgebra of $sl(2,1)$ of the) standard basis: $\epsilon_i(E_{jk}) = \delta_{i,j}\delta_{i,k}$, and $\lambda_1 = \epsilon_3$. We will denote the set of simple roots by Γ .

There are six possible Dynkin diagrams:

- (1) $\Gamma(D_1) = \{\epsilon_1 - \epsilon_2, \epsilon_2 - \lambda_1\}$. We will set $\alpha_1 = \epsilon_1 - \epsilon_2$ and $\alpha_2 = \epsilon_2 - \lambda_1$. α_1 is even, while α_2 is odd. The third positive root in this case is $\alpha_1 + \alpha_2$ which is odd.
- (2) $\Gamma(D_2) = \{\epsilon_1 - \lambda_1, \lambda_1 - \epsilon_2\}$. Note that these two roots are actually $\alpha_1 + \alpha_2$ and $-\alpha_2$, and it is easy to see that D_2 is obtained from D_1 via the odd reflection σ_{α_2} associated to the root α_2 . We can obtain D_1 back from D_2 by $\sigma_{-\alpha_2}$. Note also that the third positive root in this case will be α_1 which is even.
- (3) $\Gamma(D_3) = \{\lambda_1 - \epsilon_1, \epsilon_1 - \epsilon_2\} = \{-\alpha_1 - \alpha_2, \alpha_1\}$. We note that D_3 is obtained from D_2 via the odd reflection $\sigma_{\alpha_1 + \alpha_2}$. Applying $\sigma_{-\alpha_1 - \alpha_2}$ to D_3 will return D_2 as expected. Note also that the third positive root will be $-\alpha_2$ which is odd.
- (4) $\Gamma(D_4) = -\Gamma(D_1) = \{-\alpha_1, -\alpha_2\}$. The third positive root in this case will be $-\alpha_1 - \alpha_2$ which is odd.
- (5) $\Gamma(D_5) = -\Gamma(D_2) = \{-\alpha_1 - \alpha_2, \alpha_2\}$. The third positive root in this case will be $-\alpha_1$ which is even. Note that D_5 is obtained from D_4 via $\sigma_{-\alpha_2}$, and that applying σ_{α_2} to D_5 will yield D_4 as expected.
- (6) $\Gamma(D_6) = -\Gamma(D_3) = \{\alpha_1 + \alpha_2, -\alpha_1\}$. The third positive root will be α_2 which is odd. The odd reflections $\sigma_{-\alpha_1 - \alpha_2}$ and $\sigma_{\alpha_1 + \alpha_2}$ will map D_5 and D_6 into one another.

Hence, up to sign, there are three Dynkin diagrams, and these can be obtained from one another via a chain of odd reflections (which change the signs of some of the odd roots but a positive even root stays positive). Also note that except for the two Dynkin diagrams D_2 and D_5 , the diagrams consist of one white circle and one gray circle (standing for two roots of different parities), and so these diagrams will not allow any nontrivial admissible triples. Therefore the construction of our theorem will only yield standard r-matrices for these cases. In D_2 and D_5 , both simple roots are odd, and we can actually consider a nontrivial admissible triple in these cases. Therefore the theorem will give us one standard r-matrix and two nonstandard r-matrices for each of the diagrams D_2 and D_5 .

7.2. The Standard r-matrices. By construction, given $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ satisfying $r_0 + r_0^{21} = \Omega_0$ ¹⁰, the standard r-matrix for a fixed Dynkin diagram is

$$r = r_0 + \sum_{\alpha > 0} e_{-\alpha} \otimes e_{\alpha}.$$

So fixing r_0 we write down the standard r-matrices for the above diagrams:

- (1) For D_1 the positive roots are α_1, α_2 and $\alpha_1 + \alpha_2$. We let:

$$e_{\alpha_1} = E_{12}, \quad e_{\alpha_2} = E_{23}, \quad e_{\alpha_1 + \alpha_2} = E_{13}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{-\alpha_1} = E_{21}, \quad e_{-\alpha_2} = E_{32}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Therefore we get:

$$r_{st}(D_1) = r_0 + (E_{21} \otimes E_{12}) + (E_{32} \otimes E_{23}) + (E_{31} \otimes E_{13})$$

¹⁰ Ω_0 is the Cartan part of the quadratic Casimir element Ω of $\mathfrak{g}$.

- (2) For D_2 the positive roots are α_1 , $-\alpha_2$ and $\alpha_1 + \alpha_2$. We let:

$$e_{\alpha_1} = E_{12}, \quad e_{-\alpha_2} = E_{32}, \quad e_{\alpha_1 + \alpha_2} = E_{13}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{-\alpha_1} = E_{21}, \quad e_{\alpha_2} = -E_{23}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Therefore we get:

$$r_{st}(D_2) = r_0 + (E_{21} \otimes E_{12}) - (E_{23} \otimes E_{32}) + (E_{31} \otimes E_{13})$$

- (3) For D_3 the positive roots are α_1 , $-\alpha_2$ and $-\alpha_1 - \alpha_2$. We let:

$$e_{\alpha_1} = E_{12}, \quad e_{-\alpha_2} = E_{32}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{-\alpha_1} = E_{21}, \quad e_{\alpha_2} = -E_{23}, \quad e_{\alpha_1 + \alpha_2} = -E_{13}$$

Therefore we get:

$$r_{st}(D_3) = r_0 + (E_{21} \otimes E_{12}) - (E_{23} \otimes E_{32}) - (E_{13} \otimes E_{31})$$

- (4) For D_4 the positive roots are $-\alpha_1$, $-\alpha_2$ and $-\alpha_1 - \alpha_2$. We let:

$$e_{-\alpha_1} = E_{21}, \quad e_{-\alpha_2} = E_{32}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{\alpha_1} = E_{12}, \quad e_{\alpha_2} = -E_{23}, \quad e_{\alpha_1 + \alpha_2} = -E_{13}$$

Therefore we get:

$$r_{st}(D_4) = r_0 + (E_{12} \otimes E_{21}) - (E_{23} \otimes E_{32}) - (E_{13} \otimes E_{31})$$

- (5) For D_5 the positive roots are $-\alpha_1$, α_2 and $-\alpha_1 - \alpha_2$. We let:

$$e_{-\alpha_1} = E_{21}, \quad e_{\alpha_2} = E_{23}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{\alpha_1} = E_{12}, \quad e_{-\alpha_2} = E_{32}, \quad e_{\alpha_1 + \alpha_2} = -E_{13}$$

Therefore we get:

$$r_{st}(D_5) = r_0 + (E_{12} \otimes E_{21}) + (E_{32} \otimes E_{23}) - (E_{13} \otimes E_{31})$$

- (6) For D_6 the positive roots are $-\alpha_1$, α_2 and $\alpha_1 + \alpha_2$. We let:

$$e_{-\alpha_1} = E_{21}, \quad e_{\alpha_2} = E_{23}, \quad e_{\alpha_1 + \alpha_2} = E_{13}$$

Then we choose $e_{-\alpha}$ by $(e_{\alpha}, e_{-\alpha}) = 1$:

$$e_{\alpha_1} = E_{12}, \quad e_{-\alpha_2} = E_{32}, \quad e_{-\alpha_1 - \alpha_2} = E_{31}$$

Therefore we get:

$$r_{st}(D_6) = r_0 + (E_{12} \otimes E_{21}) + (E_{32} \otimes E_{23}) + (E_{31} \otimes E_{13})$$

Summarizing we have:

- (1) $r_{st}(D_1) = r_0 + (E_{21} \otimes E_{12}) + (E_{32} \otimes E_{23}) + (E_{31} \otimes E_{13})$
- (2) $r_{st}(D_2) = r_0 + (E_{21} \otimes E_{12}) - (E_{23} \otimes E_{32}) + (E_{31} \otimes E_{13})$
- (3) $r_{st}(D_3) = r_0 + (E_{21} \otimes E_{12}) - (E_{23} \otimes E_{32}) - (E_{13} \otimes E_{31})$
- (4) $r_{st}(D_4) = r_0 + (E_{12} \otimes E_{21}) - (E_{23} \otimes E_{32}) - (E_{13} \otimes E_{31})$
- (5) $r_{st}(D_5) = r_0 + (E_{12} \otimes E_{21}) + (E_{32} \otimes E_{23}) - (E_{13} \otimes E_{31})$
- (6) $r_{st}(D_6) = r_0 + (E_{12} \otimes E_{21}) + (E_{32} \otimes E_{23}) + (E_{31} \otimes E_{13})$

We note that the first three (and similarly the last three) are connected via odd reflections which correspond to the odd reflections that connect the associated Dynkin diagrams. The even reflection which changes the signs of the even roots will connect the first three to the last three. Hence all these r-matrices are related to one another via (even or odd) reflections.

7.3. Constructing Nonstandard r-matrices. For any given admissible triple $(\Gamma_1, \Gamma_2, \tau)$, we define a partial order on the set of positive roots, and then according to this setup, the r-matrix we obtain from our theorem is of the form:

$$r = r_0 + \sum_{\alpha > 0} e_{-\alpha} \otimes e_{\alpha} + \sum_{\alpha, \beta > 0, \alpha < \beta} (e_{-\alpha} \otimes e_{\beta} - (-1)^{|\alpha|} e_{\beta} \otimes e_{-\alpha})$$

where the particular $r_0 \in \mathfrak{h} \otimes \mathfrak{h}$ has to satisfy $(\tau(\alpha) \otimes 1)(r_0) + (1 \otimes \alpha)(r_0) = 0$ for all $\alpha \in \Gamma_1$. [Of course we still assume $r_0 + r_0^{21} = \Omega_0$].

In our case then, the nonstandard r-matrices come from the two Dynkin diagrams D_2 and D_5 :

For D_2 let $\Gamma_1 = \{\alpha_1 + \alpha_2\}$ and $\Gamma_2 = \{-\alpha_2\}$. Define $\tau(\alpha_1 + \alpha_2) = -\alpha_2$. Then the partial order on positive roots will be: $\alpha_1 + \alpha_2 < -\alpha_2$. If r_0 is given as above (i.e. $(-\alpha_2 \otimes 1)(r_0) + (1 \otimes (\alpha_1 + \alpha_2))(r_0) = 0$), then the associated r-matrix will be:

$$\begin{aligned} r_{ns_1} &= r_0 + (E_{21} \otimes E_{12}) - (E_{23} \otimes E_{32}) + (E_{31} \otimes E_{13}) \\ &+ ((E_{31} \otimes E_{32}) + (E_{32} \otimes E_{31})) \end{aligned}$$

The first few terms will actually make up $r_{st}(D_2)$ for the chosen r_0 , so we can rewrite the above as:

$$r_{ns_1} = r_{st}(D_2) + (E_{31} \otimes E_{32}) + (E_{32} \otimes E_{31})$$

For D_5 let $\Gamma_1 = \{\alpha_2\}$ and $\Gamma_2 = \{-\alpha_1 - \alpha_2\}$. Define $\tau(\alpha_2) = -\alpha_1 - \alpha_2$. Then the partial order on positive roots will be: $\alpha_2 < -\alpha_1 - \alpha_2$. If r_0 is given as above (i.e. $((-\alpha_1 - \alpha_2) \otimes 1)(r_0) + (1 \otimes \alpha_2)(r_0) = 0$), then the associated r-matrix will be:

$$\begin{aligned} r_{ns_2} &= r_0 + (E_{12} \otimes E_{21}) + (E_{32} \otimes E_{23}) - (E_{13} \otimes E_{31}) \\ &+ ((E_{32} \otimes E_{31}) + (E_{31} \otimes E_{32})) \end{aligned}$$

The first few terms will actually make up $r_{st}(D_5)$ for the chosen r_0 , so we can rewrite the above as:

$$r_{ns_2} = r_{st}(D_5) + (E_{32} \otimes E_{31}) + (E_{31} \otimes E_{32})$$

Note that if for D_2 we let $\tau(-\alpha_2) = \alpha_1 + \alpha_2$, then we would have: $-\alpha_2 < \alpha_1 + \alpha_2$, and we would get

$$r_{ns_3} = r_{st}(D_2) + (-E_{23} \otimes E_{13}) + (-E_{13} \otimes E_{23});$$

and if for D_5 we let $\tau(-\alpha_1 - \alpha_2) = \alpha_2$, then the order would be: $-\alpha_1 - \alpha_2 < \alpha_2$, and we would get

$$r_{ns_4} = r_{st}(D_5) + (-E_{13} \otimes E_{23}) + (-E_{23} \otimes E_{13}).$$

Hence the nonstandard r-matrices that can be constructed using our theorem are:

- (1) $r_{ns_1} = r_{st}(D_2) + (E_{31} \otimes E_{32}) + (E_{32} \otimes E_{31});$
- (2) $r_{ns_2} = r_{st}(D_5) + (E_{31} \otimes E_{32}) + (E_{32} \otimes E_{31});$
- (3) $r_{ns_3} = r_{st}(D_2) + (-E_{13} \otimes E_{23}) + (-E_{23} \otimes E_{13});$
- (4) $r_{ns_4} = r_{st}(D_5) + (-E_{13} \otimes E_{23}) + (-E_{23} \otimes E_{13}).$

8. CONCLUSION

In the Lie algebra case, the main classification theorem comes in two parts. The constructive part that gives an r -matrix for a given admissible triple is accompanied with the assertion that any given r -matrix that satisfies $r + r^{21} = \Omega$ can be obtained by the same construction for a suitable choice of an admissible triple. We would like to prove such an assertion for Lie superalgebras, or come up with a counterexample.

We consider once again the simple Lie superalgebra $sl(2, 1)$. We define:

$$\begin{aligned} f(E_{11} + E_{33}) &= 0 & f(E_{22} + E_{33}) &= E_{22} + E_{33} \\ f(E_{21}) &= 0 & f(E_{12}) &= E_{12} \\ f(E_{23}) &= 0 & f(E_{13}) &= E_{13} \\ f(E_{31}) &= -E_{13} & f(E_{32}) &= E_{23} + E_{32} \end{aligned}$$

and extend f to a linear map on $\mathfrak{g}$. We can easily check that this function satisfies

$$(f - 1)[f(x), f(y)] = f([(f - 1)(x), (f - 1)(y)]),$$

which is equivalent to the associated 2-tensor being an r -matrix.¹¹

We write the quadratic Casimir element:

$$\Omega = \Omega_0 + (E_{12} \otimes E_{21} + E_{21} \otimes E_{12}) + (-E_{13} \otimes E_{31} + E_{31} \otimes E_{13}) + (-E_{23} \otimes E_{32} + E_{32} \otimes E_{23})$$

where $\Omega_0 = (E_{11} + E_{33}) \otimes (-E_{22} - E_{33}) + (-E_{22} - E_{33}) \otimes (E_{11} + E_{33})$. Then if we define $r(f)$ to be the 2-tensor $(f \otimes 1)\Omega$, we get:

$$r(f) = r_0 + E_{12} \otimes E_{21} - E_{13} \otimes E_{31} + E_{32} \otimes E_{23} - E_{13} \otimes E_{13} + E_{23} \otimes E_{23}$$

where $r_0 = (-E_{22} - E_{33}) \otimes (E_{11} + E_{33})$. It is clear that $r(f)$ satisfies Equation 3.

This r -matrix is not among those constructed using Theorem 2. In fact we can prove that the two subsuperalgebras $Im(f)$ and $Im(f - 1)$ will never be simultaneously isomorphic to root subsuperalgebras. The corresponding subsuperalgebras for functions constructed by the theorem will always be root subsuperalgebras. Thus the Belavin-Drinfeld type data we used is not enough to classify all solutions to the system of Equations 3 and 4. We hope to address this problem in a separate paper.

REFERENCES

- [1] Belavin A.A., Drinfeld V.G.; "Solutions of the Classical Yang-Baxter Equation and Simple Lie Algebras", *Funct. Anal. and its Appl.* **16** (1982) pp.159-180.
- [2] Belavin A.A., Drinfeld V.G.; "Triangle Equation and Simple Lie Algebras", *Soviet Scientific Reviews, Sect. C*, **4** (1984) pp.93-165.
- [3] Etingof P., Schiffmann O.; *Lectures on Quantum Groups*, International Press, 1998
- [4] Kac, V. G.; "Lie superalgebras", *Advances in Mathematics*, **26** (1977) pp. 8-96.
- [5] Leites, D., Serganova, V.; "Defining Relations for Classical Lie superalgebras I - Superalgebras with Cartan matrix or Dynkin-type Diagram", *Topological and Geometrical Methods in Field Theory (Turku 1991)*, World Sci. Publishing, 1992, pp. 194-201.
- [6] Shchepochkina, I. M.; "Maximal solvable Subalgebras of Lie superalgebras $gl(m|n)$ and $sl(m|n)$ ", translation in *Functional Anal. Appl.* **28** (1994), no. 2, pp. 147-149

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA

E-mail address: gizem@math.berkeley.edu

¹¹Equivalently we can see that the Cayley transform Θ is an isomorphism: Θ maps $\overline{(f - 1)(x)}$ to $\overline{f(x)}$. In $C_1/C_1^\perp$ the only nontrivial coset is $\overline{E_{13} + E_{31}}$, and in $C_2/C_2^\perp$ it is $\overline{E_{23} + E_{32}}$. So we can choose $x = E_{32} - E_{13} - E_{31}$.